



AUSTRALIAN INFLUENZA SURVEILLANCE SUMMARY REPORT

No.19, 2009, REPORTING PERIOD:
12 September 2009 – 18 September 2009

Key Indicators

The counting of every case of pandemic influenza is no longer feasible in the PROTECT phase. Influenza activity and severity in community is instead monitored by the surveillance systems listed below.

Is the situation changing?	Indicated by: laboratory confirmed cases reported to NetEpi/NNDSS; GP Sentinel ILI Surveillance; and ED presentations of ILI at sentinel hospitals (NSW and WA). Laboratory data are used to determine the proportion of pandemic (H1N1) 2009 circulating in the community.
How severe is the disease, and is severity changing?	Indicated by: number of hospitalisations, ICU admissions and deaths from sentinel hospital surveillance; emergence of more severe clinical picture in hospitalised cases and ICU admissions.
Is the virus changing?	Indicated by: emergence of drug resistance or gene drift/shift from laboratory surveillance.
What is ahead?	Forward projections of cases, morbidity and mortality.

Key Points

Is the situation changing?

- As of 18 September 2009 there were 36,209 confirmed cases of pandemic (H1N1) 2009 in Australia. The number of cases reported represents a small proportion of pandemic (H1N1) 2009 cases which have occurred in the community. **The number of new confirmed pandemic (H1N1) 2009 cases and hospitalisations has decreased nationally with a number of jurisdictions reporting no new notifications in the last week, indicating that the first wave of the pandemic has subsided.**
- Overall, current national influenza notifications continue to decrease. Most jurisdictions have reported that pandemic (H1N1) 2009 activity has peaked and is decreasing.
 - ILI presentations to General Practitioners have remained steady or decreased in most states and territories with NSW reporting an increase. At a national level, rates are below levels seen at the same time in 2007 and 2008.
 - ILI presentations to emergency departments decreased across all reporting systems this reporting period.
 - Absenteeism rates decreased in the last week to levels similar to those seen at the same time in 2007.
- Type A influenza is the predominant seasonal influenza type reported by all jurisdictions. The pandemic strain has almost replaced the current seasonal H1N1 virus. The proportion of influenza positive tests that were pandemic (H1N1) 2009 increased slightly for this reporting period, with an average proportion of 93%. Of the seasonal influenza A notifications, A/H3N2 is the predominant subtype reported by most jurisdictions.

How severe is the disease?

- The number of people with pandemic (H1N1) 2009 requiring hospitalisation continues to decrease. **Seven jurisdictions have reported no new hospitalisations in the week ending 18 September 2009.** In total, 4,720 people have been hospitalised, with 13 % admitted to Intensive Care Units. The overall hospitalisation rate is 21 per 100,000 population with the highest rates in children aged less than 5 years of age (60.9 for males and 47.7 for females per 100,000 population).

- Due to the presence of underlying chronic disease, some of which is undiagnosed, and their higher level of social disadvantage, Indigenous Australians are vulnerable to complications from the pandemic (H1N1) 2009 virus. **Indigenous Australians are approximately 10 times more likely than non-Indigenous Australians to be hospitalised for pandemic (H1N1) 2009.** Of the hospitalisations for which Indigenous status is known, 773 (20%) have been Indigenous Australians.
- **Pregnant women represent 32% of all hospitalisations for pandemic (H1N1) 2009 of women aged between 20-39 years**, reinforcing the fact that pregnancy, particularly in the second and third trimesters, is an important risk factor for hospitalisation with pandemic (H1N1) 2009.
- Since reporting began, the Australian Paediatric Surveillance Unit (APSU) has reported a total of 124 notifications of children hospitalised with severe complications of influenza. Sixty percent had no underlying medical conditions.
- As of 18 September 2009, the number of deaths associated with pandemic (H1N1) 2009 was 172. Of these deaths, 3 were pregnant women and 23 (13%) were Indigenous.
- The median age of confirmed cases that died is 51 years (range 2-86 years of age), compared to the median age for deaths from seasonal flu from 2001 to 2006 which is 83 years.
- Reports from the Australian jurisdictions indicate that most of the deaths had underlying medical conditions including cancer, diabetes mellitus and morbid obesity.

Is the virus changing?

- In Australia, the WHO Collaborating Centre for Reference and Research on Influenza has tested 250 pandemic (H1N1) 2009 viral isolates by NA enzyme inhibition assay and 123 clinical specimens were tested for the H275Y mutation (known to confer resistance to oseltamivir). Of these, two clinical specimens were positive for the H275Y mutation and one was also resistant to oseltamivir when tested by NA enzyme inhibition assay.
- To date, the WHO has received formal notification of 26 cases of oseltamivir resistance pandemic (H1N1) 2009 viruses worldwide.

What is ahead?

- With a 20% clinical attack rate and no intervention; it has been estimated that by the end of winter 1 in 5 Australian (4.3 million) could become infected with the pandemic virus, leading to 80,000 hospitalisations and 6,000 deaths. Currently the number of hospitalisations and deaths are tracking below these estimations.


1. Current influenza activity in Australia – Is the situation changing?

Notifications of confirmed pandemic (H1N1) 2009 and seasonal influenza

The number of new notifications of laboratory confirmed pandemic (H1N1) 2009 this reporting period was 181. As of 18 September 2009 there were 36,209 confirmed cases of pandemic (H1N1) 2009 in Australia, including 172 associated deaths. The number of new confirmed pandemic (H1N1) 2009 cases and hospitalisations has decreased nationally with a number of jurisdictions reporting no new notifications in the last week, indicating that the first wave of the pandemic has subsided. The number of cases reported represents only a small proportion of pandemic (H1N1) 2009 circulating in the community.

The national epidemic curve shows the jurisdictional distribution of confirmed cases of pandemic (H1N1) 2009 over time in Australia (Figure 1).

Figure 1. Laboratory confirmed cases of pandemic (H1N1) 2009 in Australia, to 18 September 2009 by jurisdiction




Source: NetEPI database

As Figure 2 shows, influenza activity in 2009 started earlier than in 2008 and there was a rapid increase in the number of confirmed influenza cases (both seasonal and pandemic (H1N1) 2009) from week 21 (starting 16 May 2009). The high number of confirmed notifications of seasonal influenza seen during May and June are most likely due to the increase in testing for pandemic (H1N1) 2009. Overall, numbers of laboratory confirmed notifications of influenza have been decreasing in the past few weeks.

Laboratory confirmed notifications of influenza for this reporting week are 0.9 times the 5 year weekly mean and the notifications year to date are 9.6 times the 5 year, year to date mean.

Figure 2. Influenza activity in Australia, by reporting week, years 2007, 2008 and 2009*



* Data on pandemic (H1N1) 2009 cases is extracted from NetEPI; data on seasonal influenza is extracted from the NNDSS. Sources: NNDSS and NetEPI databases

On 17 June 2009 Australia commenced the transition to a new response phase called PROTECT, in which laboratory testing is directed towards people with moderate or severe illness; those more vulnerable to severe illness; and those in institutional settings. This means that the number of confirmed cases does not reflect how many people in the community have acquired pandemic (H1N1) 2009 infection.

As the counting of every case is no longer feasible in the PROTECT phase, influenza activity, including Influenza Like Illness (ILI) activity in the community, is instead monitored by surveillance systems including:

- GP Sentinel ILI surveillance;
- Emergency Department presentations of ILI at sentinel hospitals (NSW and WA); and
- Absenteeism rates.

Laboratory data are used to determine the proportion of pandemic (H1N1) 2009 circulating in the community.


Influenza Like Illness activity in Australia

Sentinel General Practice

ILI presentations to General Practitioners continue to decrease and are lower than 2007 and 2008 rates nationally. In most jurisdictions ILI data have decreased or plateaued in this period with NSW reporting an increase in activity.

Combined data available from the Australian Sentinel Practices Research Network (ASPREN), the Northern Territory GP surveillance system and VIDRL, up until 13 September 2009, show that nationally, influenza like illness (ILI) consultation rates remained stable this reporting period and are below levels seen in 2007 and 2008 (Figure 3). In the last week, the presentation rate to sentinel GPs in Australia was approximately 11 cases per 1,000 patients seen.

Figure 3. Rate of ILI reported from GP ILI surveillance systems from 2007 to 13 September 2009 by week*




* Delays in the reporting of data may cause data to change retrospectively. As data from the NT and the VIDRL surveillance systems are combined with ASPREN data, rates may not be directly comparable across 2007, 2008 and 2009.

SOURCE: ASPREN, NT, VIDRL








Further analysis of the ILI data showed levels in most jurisdictions decreased or plateaued in this period with NSW reporting an increase in activity. Victoria, Tasmania, the Australian Capital Territory and the Northern Territory are at background levels (Figure 4).

Care should be taken when interpreting Figure 4 graphs due to lags in reporting in some instances and small numbers being reported from jurisdictions. The last data point may be modified in future reports.

Figure 4. Rate of ILI reported from ASPREN, VIDRL and NT by State from January 2009 to 13 September 2009 by week



SOURCE: ASPREN (VIC) & VIDRL




SOURCE: NT Surveillance

Emergency departments

Trends in ILI presentations to EDs decreased across all reporting systems this reporting period.

The number of ILI presentations reported in Western Australian EDs has dropped in the week ending 13 September 2009 with levels similar to those seen at the same time in 2007 and 2008 (Figure 5). The proportion of ILI presentations admitted to hospital remained stable at 5.2%.

Figure 5. Number of Emergency Department presentations due to ILI in Western Australia from 1 January 2007 to 13 September 2009 by week




In early July 2007 (week 26), several deaths associated with influenza infection were reported in children from Western Australia. The public response to these deaths could account for the sudden increase in ILI presentations to Perth EDs in 2007.

SOURCE: WA 'Virus Watch' Report

In the week ending 11 September 2009, ILI presentations to New South Wales EDs decreased to low levels (rate 2.2 per 1,000 presentations) (Figure 6). Presentations were mainly for mild illnesses and 10% of presentations with ILI were admitted, which is slightly above levels reported previously.

Figure 6. Rate of ILI diagnosed in people presenting to selected Emergency Departments, NSW 1 January 2005 to 11 September 2009 by month*



* Emergency department data are preliminary and may be updated in later weeks.


SOURCE: NSW HEALTH 'NSW Influenza Surveillance Report'

ILI presentations to South Australian EDs remained steady with 32 presentations this reporting period compared with 33 presentations in the previous week. The number of admissions remained stable at 3.¹

Absenteeism

Absenteeism rates have dropped in the last week to levels similar to those seen in 2007 (Figure 7).

Figure 7. Rates of absenteeism of greater than 3 days absent, National employer, 1 January 2007 to 9 September 2009, by week.



SOURCE: Absenteeism data

**Laboratory surveillance:
How much ILI and influenza is due to pandemic (H1N1) 2009?**

As shown in Table 1 below, of those tests that were positive for Influenza A, the proportion of tests that were pandemic (H1N1) 2009 remained stable in most jurisdictions with a mean of 88%.

The proportion of pandemic (H1N1) 2009 to seasonal influenza varies across jurisdictions. This proportion is used as an indicator to help determine if a person has influenza, then how likely it is to be pandemic (H1N1) 2009. The proportion of pandemic (H1N1) 2009 to seasonal influenza as reported by the jurisdictions is shown in Table 1. The proportion of confirmed influenza in Australia which was pandemic (H1N1) 2009 across all reporting systems increased slightly to 93%.

Over the last two weeks, for the days on which surveillance testing was conducted, ASPREN GPs reported 32 people presenting with ILI. Of these, 63% (20/32) were tested for influenza and none tested positive for influenza.

Table 1. Laboratory tests that tested positive for influenza A and pandemic (H1N1) 2009

	ASPREN – national	NSW Report ^	VIC NIC	WA NIC
Latest report				
Number of specimens tested	20	756 (at 18/9)	154 (at 18/9)	403 (at 18/9)
% tested which were Influenza A	0% (0)	2% (13)	1% (2)	7% (28)
% tested which were pandemic (H1N1) 2009	0% (0)	77% (10)	100% (2)	100% (28)
Previous report				
Number of specimens tested	20	795 (at 11/9)	43 (at 11/9)	623 (at 11/9)
% tested which were Influenza A	20% (4)	3% (23)	9% (4)	10% (60)
% tested which were pandemic (H1N1) 2009	100% (4)	87% (20)	100% (4)	95% (57)

*ASPREN tests are collected every Tuesday. Results are reported for a rolling fortnight as data changes retrospectively.

^NSW Influenza Report available from: <http://www.emergency.health.nsw.gov.au/swineflu/index.asp>

The proportion of pandemic (H1N1) 2009 compared with seasonal influenza varies across different countries in both the Northern and Southern Hemispheres. Argentina reported that 93% of the respiratory viruses circulating in those aged over 5 years is due to pandemic (H1N1) 2009 and influenza A untyped², while Canada reported that pandemic (H1N1) 2009 represents 99% of all influenzas³, and the US is reporting that it represents 99% of the total circulating influenza viruses.⁴ In New Zealand, pandemic (H1N1) 2009 represents 100% of influenza viruses reported from sentinel surveillance and 50% of influenza viruses reported in non-sentinel surveillance.⁵

2. How severe is the disease, and is severity changing?

Overview of pandemic (H1N1) 2009 severity

Table 2 provides a summary of measures that indicate the severity of pandemic (H1N1) 2009 since the beginning of the outbreak and up to 18 September 2009. Of particular note is the increasing median age as the severity of the disease progresses: 21 years for all confirmed cases; 31 years for hospitalised cases; 46 years for ICU cases; and 51 years for cases who have died.

Table 2. Summary of severity indicators of pandemic (H1N1) in Australia#

	Confirmed pandemic (H1N1) 2009 cases	Hospitalised cases	ICU cases	Deaths
Total number	36,209	13% (4,720/36,209)	13% (615/4,720)	172
% Females	51 (18,399/36,080)	50% (2,395/4,720)	52% (324/615)	44% (79/172)
Median age (years)	21	31	46	51
Indigenous people	10% (3,351/33,352)	20% (773/3,801)	20% (91/466)	13% (23/172)
% Pregnant	n/a	11% (262/2,395)	7% (24/324)	2% (3/172)

#Data are extracted from a number of sources depending on the availability of information. Figures used in the analysis have been provided in parentheses.

Table 3. Summary of hospitalisations and deaths associated with pandemic (H1N1) 2009 in Australia, by Jurisdiction

	ACT	NSW	NT	QLD	SA	TAS	VIC	WA	AUS
Total pandemic (H1N1) 2009 hospitalisations	43	1239	359	1228	470	106	418	857	4,720
Percentage of national pandemic (H1N1) 2009 hospitalisations	0.9%	26.3%	7.6%	26.0%	10.0%	2.2%	8.9%	18.2%	100%
Crude rate per 100,000	12.5	17.8	163.2	28.7	29.3	21.3	7.9	40.0	22.1
Total pandemic (H1N1) 2009 deaths	2	48	7	38	21	6	24	26	172
Percentage of national pandemic (H1N1) 2009 deaths	1.2%	27.9%	4.1%	22.1%	12.2%	3.5%	14.0%	15.1%	100%
Crude rate per 100,000	0.6	0.7	3.2	0.9	1.3	1.2	0.5	1.2	0.8

The jurisdictions report directly to the National Incident Room (Commonwealth Department of Health and Ageing) on hospitalisations and deaths associated with the pandemic (H1N1) 2009 virus.

Source: NetEpi

Pandemic morbidity (hospitalisations)


Hospitalisations of Pandemic (H1N1) 2009 confirmed cases

As of 18 September 2009, jurisdictions have reported that 4,720 confirmed cases of pandemic (H1N1) 2009 have been hospitalised (this figure includes people with pandemic (H1N1) 2009 who are hospitalised for associated conditions). The number of cases per day requiring hospitalisation has decreased since late August. **Seven jurisdictions have reported no new hospitalisations in the week ending 18 September 2009** (Figure 8).

Indigenous Australians are approximately 10 times more likely than non-Indigenous Australians to be hospitalised for pandemic (H1N1) 2009. Since the beginning of the outbreak Australian jurisdictions have reported that 773 (20%) of the 3,801 cases hospitalised for which there was information on Indigenous Status were Aboriginal and/or Torres Strait Islander.

For comparative purposes, for the period 2000-01 to 2006-07, an average of 1,925 people with influenza are admitted to hospital each year. For all influenzas^a and pneumonias^b, for the same period, an average of 73,271 people were admitted to hospital.⁶

Figure 8. Weekly numbers of hospital admissions of confirmed cases, to 18 September 2009, Australia



Source: NETEPI database


Age and sex distribution of hospitalised confirmed cases

There have been 4,720 confirmed cases hospitalised since the beginning of the outbreak. Of these cases, the overall hospitalisation rate is 21.0 per 100,000 population, with the highest rates in children aged less than 5 years of age (60.9 for males and 47.7 for females per 100,000 population). The median age of hospitalised cases is 31 years (range 0-98 years). Figure 9 illustrates that the age distribution of hospitalised cases of pandemic (H1N1) 2009 is different to previous influenza seasons. In comparison with the 2004-2007 influenza seasons, young children aged less than 5 years of age continue to be hospitalised at a higher rate than other age groups (males in particular) but for pandemic (H1N1) 2009 there is a peak in the 50 to 60 years age group and a marked decrease in those aged more than 75 years.

^a ICD10-AM codes J10-J11

^b ICD10-AM codes J12-J18

Figure 9. Age specific rates of hospitalised confirmed cases of pandemic (H1N1) 2009 to 18 September 2009, compared with average annual age specific rates of hospitalisations from seasonal influenza 2004-05 to 2006-07*, Australia




*The rates for pandemic (H1N1) 2009 are from 15 June to 21 August 2009 whereas the rates for seasonal influenza are averaged annual rates (i.e. for a full influenza season).

Source: NETEPI database

Information on length of stay is available for 68% (3,220) of the 4,720 hospitalised cases for which there is further information. The median length of stay in hospital is 3 days (range 1-98 days). Approximately 19% of all hospitalised patients stayed in hospital for more than 7 days (Figure 10). A breakdown by age group shows that children aged less than 5 years, although more likely to be hospitalised, tend to be hospitalised for shorter periods than older children and adults. Only 2% of children aged less than 5 years remain in hospital for longer than 7 days compared to 13% for those in the age group 30 years and over.

Figure 10. Hospitalised confirmed cases of pandemic (H1N1) 2009, by length of hospital stay and age group, to 18 September 2009, Australia




Source: NETEPI database

Pregnancy as a risk factor for hospitalisation with pandemic (H1N1) 2009

A total of 262 (6%) of the 4,720 hospitalised confirmed cases for whom further information was known were pregnant women. Pregnant women represent 32% of all hospitalisations for pandemic (H1N1) 2009 of women aged between 20-39 years. Information on gestation is available for 68 of the 262 cases. Approximately 6% (4) were in their 1st trimester (weeks 1-12); 25% (17) were in their 2nd trimester (weeks 13-26); and 69% (47) were in their 3rd trimester (weeks 27-41) (Figure 11). A total of 39 pregnant women were admitted to ICU: 1 was in her first trimester, 4 were in their 2nd trimester and 9 were in the 3rd trimester. No information on gestation was available for 25 women. Pregnant women stayed an average of 15 days in hospital (range 1-63 days). Three pregnant women are known to have died.

Figure 11. Hospitalised confirmed cases of pandemic (H1N1) 2009 in pregnant women by weeks of gestation, to 18 September 2009, Australia




Source: NETEPI database

Paediatric hospital admissions

Since reporting began in 2009, 124 children have been reported as hospitalised with complications from influenza by the Australian Paediatric Surveillance Unit (APSU). Admission data has been provided for 85 cases (Figure 12).

Of the 89 cases, for which data are available, the average age of children admitted to hospital is four years and five months, with an age range from one month to 16 years. Complications were mostly for pneumonia and encephalitis. Thirty-three of the 82 (40%) cases for which data is available had underlying conditions.

Figure 12. Number of paediatric hospital admissions APSU, 11 March 2009 to 1 September 2009, by week of admission.



SOURCE: APSU

Confirmed cases requiring intensive care

In Week 38 (week ending 18 September 2009), an average of 62 hospitalised cases required intensive care on any given day^c a decrease from 66 the previous reporting period. Thirteen percent of hospitalised cases have required admission to ICU. Twenty percent of ICU cases have been reported as Indigenous. The median age of cases in ICU is 46 years (range 0-84 years).

Snapshot on Queensland^d hospitalised confirmed cases

There have been 1,229 hospitalisations associated with pandemic (H1N1) 2009 in Queensland between 25 May 2009 and 18 September 2009, an age standardised (non-annualised) rate of 28.7 per 100,000 population. Fifty four percent (n=659) of hospitalisations were in females, an age standardised rate of 30.8 per 100,000 population. Forty six percent (n=569) of hospitalisations were in males, an age standardised rate of 26.6 per 100,000 population.

Of these 1,229 hospitalised cases, co-morbidities were reported in 64% (n=778) of cases. Co-morbidities included chronic respiratory 53% (n=413); diabetes 15% (n=118); pregnancy 10% (n=78); chronic cardiac 12% (n=94); immuno-compromised 11% (n=86); morbid obesity 8% (n=59); and renal 5% (n=40). Information on the length of time between onset of symptoms and

^c This does not represent the number of new cases requiring admittance to an Intensive Care Unit (ICU) but is a repeated measure of the prevalence of confirmed cases in an ICU on a particular day.


^d Queensland has been able to provide greater detail on hospitalised cases than other States and Territories, enabling more in-depth analysis of hospitalised cases.

admittance to hospital were available for 895 confirmed cases. Fifty three percent (477/895) of these cases were admitted within 48 hours of onset of symptoms.

Figure 13 illustrates that the 0-4 years age group is more likely to be hospitalised but the length of stay is more likely to be less than 3 days than for other age groups. Twenty seven percent (n=275) of cases were hospitalised for less than 3 days; 48% (n=487) for 3-7 days; 11% (n=109) for 8-13 days; and 13% (n=137) for 14 or more days. These figures include cases that are still hospitalised.

Of these hospitalised cases, 185 were admitted to an ICU or special care with 54% being female. The median length of stay in ICU or special care was 9 days (range 1-66 days). The age of cases admitted to ICU ranged from 0-84 years with the highest proportion of cases in the 50-54 year age group (14%, n=26), followed by the 55-59 year age group and the 45-49 year age group, which each accounted for 13% (n=24) and 11% (n=21) of the cases respectively. Thirty eight percent (n=70) of cases in ICU had no reported co-morbidities.

Figure 13. Hospitalised confirmed cases of pandemic (H1N1) 2009, by length of hospital stay and age group, to 18 September 2009, Queensland



Source: QLD Health


Pandemic Mortality

Deaths associated with pandemic (H1N1) 2009

There were 172 deaths associated with the pandemic (H1N1) 2009 virus in Australia between 19 June 2009 and 18 September 2009.^e Of these 172 deaths, 48 occurred in New South Wales, 38 in Queensland, 26 in Western Australia, 24 in Victoria, 21 in South Australia, 7 in the Northern Territory, 6 in Tasmania, and 2 in the Australian Capital Territory. Of the 172 deaths, 23 (13%) were Indigenous.^f Reports from the jurisdictions in Australia indicate that most of the cases had underlying medical conditions including cancer, diabetes mellitus and morbid obesity.

Males were overrepresented among the deaths (52%, n=93). The median age of confirmed cases that died was 51 years (range 2-86 years). This is lower than the median age of deaths from seasonal influenza for the period 2001-2006 (83 years). The highest proportion of deaths (14%, n=25) have occurred in the 55-59 year age group. Most deaths (73%) have occurred in those aged between 35 to 79 years (Figure 14). Noting the apparent biases in these data, the pattern of deaths across age groups is very different to the age distribution of hospitalisations and confirmed cases.

Figure 14. Numbers of deaths among confirmed cases of pandemic (H1N1) 2009, by age group and sex, compared with total laboratory confirmed pandemic (H1N1) 2009 notifications by age group, to 18 September 2009, Australia



Source: NETEPI database

Deaths associated with influenza and pneumonia

There are difficulties estimating the number of deaths due to influenza in Australia. Deaths coded as being due to laboratory confirmed influenza are known to underestimate the true number. Influenza may not be listed on the death certificate if it wasn't recognised as the underlying cause. Coding of pneumonia and influenza provides an additional measure, although this will overestimate the number of deaths as it will include other causes of pneumonia.


^e For the most recent figures on hospitalisations and deaths please access the latest Situation Report at <http://www.healthemergency.gov.au/internet/healthemergency/publishing.nsf/Content/updates>

^f It is estimated that 2.4% of the total Australian population are Aboriginal and/or Torres Strait Islander.

The median number of annual deaths in Australia for the years 2001 to 2006 from influenza and pneumonia is 3,089 and for laboratory diagnosed influenza is 40. In 2007 (the latest year for which data has been released) there were 2,623 deaths with influenza and pneumonia as the underlying cause of death. In 2007, influenza and pneumonia was the 13th leading cause of death in Australia (*Source: ABS, Causes of Death 2007.*). Mortality figures are likely to be an underestimate due to inherent difficulties in assigning causes of death and therefore appropriate ICD codes. ABS mortality data are released two years in arrears.

Although mortality data from all causes are generally not available for the current year, some information on influenza and pneumonia deaths are reported by individual jurisdictions from their Births, Deaths and Marriages Registers. **In Western Australia, pneumonia and influenza deaths accounted for approximately 18% of all deaths in the last week of August, which is below levels seen at the same time in 2007 and 2008 (Figure 15).**


Figure 15: Percentage of all deaths classified as influenza and pneumonia, WA Registry of Births, Deaths and Marriages, 1 January 2008 to 30 August 2009



SOURCE: WA 'Virus Watch' Report

In NSW, death certificate data as of 28 August 2009 show that there were 75 influenza or pneumonia deaths per 1,000 deaths in NSW, which was below the expected seasonal threshold for this time of year of 154 per 1,000 (Figure 16).

Figure 16: Rates of deaths classified as influenza and pneumonia, NSW Registry of Births, Deaths and Marriages, 1 January 2004 to 28 August 2009



SOURCE: NSW Health 'Weekly Influenza Report'

3. Is the virus changing?

Laboratory Confirmed Influenza

It is not possible to determine accurately the number of notifications due to seasonal influenza. Increasingly, not all influenza viruses are subtyped and the large proportion of influenza A (13,541 notifications) reported to NNDSS could be either pandemic (H1N1) 2009 or seasonal influenza. Laboratory reports in recent weeks estimate that 93% of all influenza positive tests are due to pandemic (H1N1) 2009.

From 1 January to 18 September 2009, type A is the predominant seasonal influenza type reported by all jurisdictions. For those type A notifications in NNDSS which are not pandemic (H1N1)2009, but for which there is subtyping information, the ratio of seasonal H1N1 to H3N2 is 1:2.2.

Antigenic characteristics

WHO Collaborating Centre for Reference & Research on Influenza (WHO CC)

In 2009 up to 20 September 2009, 968 Australian influenza isolates have been subtyped by the WHO CC in Melbourne. Of these, 492 influenza isolates have been antigenically characterized.

In general, seasonal influenza A strains circulating this influenza season are the same as strains in the vaccine, with the A(H3N2) virus drifting. Influenza B strains match more closely with those in the 2009-10 Northern Hemisphere vaccine and may be drifting.

Antiviral Resistance

Pandemic (H1N1) 2009

To date, the WHO has received formal notification of 26 cases of oseltamivir resistant pandemic (H1N1) 2009 viruses worldwide. The isolates have a mutation in the neuraminidase (referred to as H275Y) that confers resistance to oseltamivir, but they remain sensitive to zanamivir. Worldwide there have been over 10,000 clinical samples and isolates of the pandemic (H1N1) 2009 tested and found to be sensitive to oseltamivir. WHO continues to monitor the situation closely.⁷

In Australia, the WHO Collaborating Centre for Reference and Research on Influenza has tested 250 pandemic (H1N1) 2009 viral isolates by NA enzyme inhibition assay and 123 clinical specimens were tested for the H275Y mutation (known to confer resistance to oseltamivir). Of these, two clinical specimens were positive for the H275Y mutation and one was also resistant to oseltamivir when tested by NA enzyme inhibition assay.

The US CDC has reported a total of 10 cases of oseltamivir resistance strains of pandemic (H1N1) 2009 in the week ending 12 September 2009. All cases except one had exposure to oseltamivir through either treatment or prophylaxis. One case is currently under investigation to determine exposure to oseltamivir.⁴ The US CDC reported on 11 September 2009 that 2 cases involved adolescent girls receiving oseltamivir as prophylaxis in a summer camp. Both girls stayed in the same cabin at the camp and specimens from both girls had the H275Y mutation, along with a second mutation (I223V) in neuraminidase. This is the first report of oseltamivir resistance in Pandemic (H1N1) 2009 cases with an epidemiologic link.⁸ All of the 670 pandemic (H1N1) viral isolates tested by the US CDC continued to be sensitive to zanamivir.⁴

In New Zealand, all of the 92 pandemic (H1N1) 2009 viruses tested up to 13 September 2009 were sensitive to oseltamivir, including one from a fatal case of a 21 year-old male. None of the 12 pandemic (H1N1) 2009 clinical specimens tested positive for the H275Y mutation which confers resistance to oseltamivir.⁵

The UK HPA has reported that 2 of 913 pandemic viruses tested in England have been confirmed to carry a mutation which confers resistance to oseltamivir, and one of these has been shown phenotypically to be resistant to the drug but remains sensitive to zanamivir.⁹

Seasonal Influenza

The last WHO report on resistance of seasonal strains to oseltamivir was released on 4 June 2009, during the Northern Hemisphere influenza season 2008-2009. This report stated that 96% of seasonal influenza A (H1N1) isolates tested from 36 countries worldwide were resistant to oseltamivir.¹⁰

In Australia (since 1 January 2009) 36 of the 37 seasonal H1N1 viruses tested were resistant to oseltamivir. All of the 40 A(H3N2) viruses and the 6 influenza B viruses tested were sensitive to both oseltamivir and zanamivir.

In New Zealand, all of the 28 seasonal A(H1N1) viruses have been tested up to 13 September 2009 for the H275Y mutation (which is known to confer resistance to oseltamivir) had the mutation.⁵

The US CDC reported in the week ending 12 September that 1,143 of the 1,148 seasonal A(H1N1) isolates tested were resistant to oseltamivir and all of the 261 Influenza A(H3N2) isolates tested were resistant to adamantanes.⁴

Data considerations

The information in this report is reliant on the surveillance sources available to the Department of Health and Ageing. As access to sources increase and improve, this report will be refined and additional information will be included.

This report aims to increase awareness of pandemic (H1N1) 2009 and seasonal influenza in Australia by providing an analysis of the various surveillance data sources throughout Australia. While every care has been taken in preparing this report, the Commonwealth does not accept liability for any injury or loss or damage arising from the use of, or reliance upon, the content of the report. Please note, the pandemic (H1N1) 2009 and seasonal influenza elements of this report are based on data available as at 18 September 2009. Delays in the reporting of data may cause data to change retrospectively. For further details about information contained in this report please contact the Influenza Team through flu@health.gov.au.

NetEpi

All jurisdictions except QLD are reporting pandemic (H1N1) 2009 cases using NetEpi, a web-based outbreak case reporting system. Data from jurisdictional systems are being imported into NetEpi by VIC, NSW, WA, TAS and SA, and the remainder are entering directly into NetEpi. QLD ceased reporting into NetEpi on 6 July 2009.

Analyses of Australian cases are based on clinical onset date, if this information is available. Where an onset date is not available, notification date has been used. Victorian cases use a calculated onset date which is the earliest available date calculated from specimen date, onset date, notification date or detection date. This assumption was made for all calculations and data on which the figures are based.

State and Territory reporting

The jurisdictions report directly to the National Incident Room, Commonwealth Department of Health and Ageing, on hospitalisations, numbers admitted to ICUs and deaths.

National Notifiable Diseases Surveillance System (NNDSS)

NNDSS comprises of notifications from jurisdictions of laboratory-confirmed influenza cases. Laboratory confirmed influenza is notifiable in all jurisdictions in Australia. Confirmed pandemic (H1N1) 2009 cases are being received from all jurisdictions through NNDSS except for Victoria and New South Wales. NSW is also unable to send seasonal influenza notifications data.

Data Analysis

Analysis of confirmed cases is done on combined NetEpi and NNDSS data. Analysis of morbidity (hospitalisations and ICU admissions) and mortality data is done on combined NetEpi and QLD hospitalisation data.

Laboratory Surveillance data

Laboratory testing data are extracted from the 'NSW Influenza Report,' 'The 2009 Victorian Influenza Vaccine Effectiveness Audit Report' (VIDRL) and the 'South Australian Seasonal Influenza Report'. These reports are provided weekly.

WHO Collaborating Centre for Reference & Research on Influenza (WHO CC)

Data are provided weekly to the Surveillance Branch from the WHO CC.

Sentinel General Practice Surveillance

The Australian Sentinel Practices Research Network (ASPREN) has Sentinel GPs who report influenza-like-illness (ILI) presentation rates in NSW, SA, ACT, VIC, QLD, TAS and WA. As jurisdictions joined ASPREN at different times and the number of GPs reporting has changed over time, the representativeness of ASPREN data in 2009 may be different from that of previous years. ASPREN data are sent to the Surveillance Branch on a weekly basis. Northern Territory GP surveillance data are sent to the Surveillance Branch on a weekly basis. VIDRL influenza surveillance data are sent to the Surveillance Branch on a weekly basis.

A new testing protocol introduced through ASPREN requires GPs to test all patients presenting with an ILI on one day of the week. These data should provide a cross section of age, sex and severity of patients who seek GP assistance for ILI. This system is in the early stages of implementation and will be further developed over coming weeks.

Sentinel Emergency Department (ED) data

WA - ED surveillance data are extracted from the 'Virus Watch' Report. This report is provided weekly. The Western Australia Influenza Surveillance Program collects data from 8 Perth Emergency Departments (EDs).

NSW - ED surveillance data are extracted from the 'NSW Influenza Surveillance Report'. This report is provided weekly. The New South Wales Influenza Surveillance Program collects data from 49 EDs across New South Wales.

SA – ED surveillance data are extracted from the 'South Australian Seasonal Influenza Report'. This report is provided weekly. The South Australian Influenza Surveillance Program collects data from 4 EDs in South Australia.

Absenteeism

A national organisation provides data on the number of employees who have been on sick leave for a continuous period of more than three days. These data are not influenza or ILI specific and absenteeism may be a result of other illnesses.

Mortality data

Mortality data are extracted from the NSW Health 'Weekly Influenza Epidemiology Report' and the WA 'Virus Watch' Report.

Paediatric hospital admissions data

Reports of ICU admissions are provided to the Surveillance Branch on a weekly basis by the Australian Paediatric Surveillance Unit. APSU conducts surveillance of severe complications of influenza in children aged 15 years and under. Surveillance began on 1 June 2009.

References

- ¹ South Australian Seasonal Influenza Report No.17. Available from: <http://www.dh.sa.gov.au/pehs/notifiable-diseases-summary/flu-resp-intro.htm>.
- ² Argentina Ministry of Health, Pandemic Situational Report. Available from: <http://www.msal.gov.ar/hm/site/default.asp>. Accessed 22 September 2009.
- ³ Canada Public Health Agency, Flu Watch: <http://www.phac-aspc.gc.ca/fluwatch/08-09/>. Accessed 2 September 2009.
- ⁴ CDC Influenza reports <http://www.cdc.gov/h1n1flu/pubs/>. Accessed 22 September 2009.
- ⁵ New Zealand Public Health Surveillance, Influenza Weekly Updates. Available at: http://www.surv.esr.cri.nz/virology/influenza_weekly_update.php. Accessed 22 September 2009.
- ⁶ Australian Institute of Health and Welfare (AIHW) National Hospital Morbidity Database. Available at: <http://www.aihw.gov.au/hospitals/datacubes/index.cfm>
- ⁷ WHO Pandemic (H1N1) 2009 update 65. Available from: <http://www.who.int/csr/disease/swineflu/updates/en/index.html>. Accessed 22 September 2009.
- ⁸ CDC Morbidity and Mortality Weekly Report <http://www.cdc.gov/mmwr/PDF/wk/mm5835.pdf>. Accessed 16 September 2009.
- ⁹ HPA Pandemic Situational Update. Available from: <http://www.hpa.org.uk/webw/HPAweb&Page&HPAwebAutoListName/Page/1240732817665?p=1240732817665>. Accessed 22 September 2009.
- ¹⁰ WHO Influenza A virus resistance to oseltamivir and other antiviral medicines, 4 June 2009. Available from: <http://www.who.int/csr/disease/influenza/en/>. Accessed 12 August 2009.